

RESEARCH

Open Access



A structural equation model linking health literacy, self efficacy and quality of life in adults with coronary heart disease

Shaoying Du¹, Zhimin Feng², Wen Wang¹, Licong Tian¹ and Yan Wang^{1*}

Abstract

Background: Cardiovascular disease is the world major cause of death. There is sufficient evidence that patients with coronary heart disease (CHD) experience poor quality of life. Health literacy and self efficacy are modifiable psychosocial factors that could affect quality of life, and these factors should be considered as targets for intervention. As the relationships among health literacy, self efficacy, and quality of life in the CHD population have not been well understood. Thus, we constructed the structure equation model in these valuables.

Methods: A cross-sectional study of a convenience sample among 200 patients with CHD were participated from outpatient clinics in three tertiary general hospitals in Baoding City in mainland China, from December 2018 to June 2019. Data regarding demographic features, health literacy, self efficacy and quality of life were assessed. A structure equation model was used to construct and validate the pathways.

Results: The mean age of the study sampled patients was 65.37 years old. The average level of health literacy, self efficacy and quality of life were 9.6 ± 3.5 , 28.8 ± 13.9 and 381.8 ± 130.1 respectively. Significant associations were observed from health literacy to quality of life, and self efficacy played a partial mediating role between health literacy and quality of life in the CHD population. Health literacy and self efficacy explained for 59.6% of the variance in quality of life.

Conclusions: Health literacy had a direct influence on quality of life, and an indirect influence on quality of life via self efficacy in the patients with CHD.

Keywords: Coronary heart disease, Health literacy, Self efficacy, Quality of life, Structural equation modelling

Background

Cardiovascular disease (CVD) is the world 1st leading cause of death in the twentieth century [1]. There are 290 million CVD patients in China, among which 11 million are coronary heart disease (CHD) patients [2]. Although the mortality have declined by the improved treatment, the CHD patients have to cope with the symptoms such

as chest pain, breathing difficulties, as well as complex treatment options for a long time, which could negatively impact the patients' satisfaction and physical and mental health. The dimensions of Quality of life (QOL) include physical and mental health. QOL is regarded as an important outcome of health care and is increasingly viewed as a comprehensive indicator of health in population health surveys and nursing interventions [3].

There is sufficient evidence that patients with CHD experience poor QOL comparing with the healthy population [4]. One research indicates that there is limited understanding of QOL and its influencing factors in

*Correspondence: chjanwy@163.com

¹ School of Nursing, Hebei University, No. 342 Yuhudong Road, Baoding City 071002, Hebei Province, People's Republic of China
Full list of author information is available at the end of the article



CHD population in Asia [5]. Understanding the influencing factors of QOL is essential to alleviate the effect of CHD on public health. It was shown that QOL could be affected by many factors, such as socio-demographic factors (i.e., age, gender, education level and social support), physical factors (i.e., health status and comorbidity), and psychological factors (i.e., stress, anxiety and depression) [6–9]. However, some of these factors are objective, and they cannot be modified [8, 9]. Moreover, health care givers maybe limited from enhancing QOL, as only a partial picture of QOL and associated factors were given [6]. Therefore, to recognize the importance of an individual's holistic perspective is needed.

HL is regarded as individuals' capacity to obtain, process, understand the health information and make appropriate healthcare decisions [10]. Therefore, HL is an important part of secondary prevention in CHD patients. Because these patients could read and understand health education materials, adhere to medication regimes, make lifestyle changes, and manage their condition effectively by HL [11], which lead them to improving the QOL. Recent report has shown that HL is now emerging as a significant determinant to improve self-rated health [12].

HL was directly associated with QOL in research probing the association in patients with ischaemic heart disease [13] and in percutaneous coronary intervention (PCI) patients at 6 months after discharge [14]. However, the findings were inconstantly observed in different studies. One study in hospitalized CHD patients found that there was no direct correlation between HL and QOL, but only indirect correlation through the mediator of SE [15]. The mixed findings showed that the relationship among HL, SE and QOL might be complicated and require further clarification.

SE refers to an individual's confidence in one's ability to complete tasks [16]. Paasche-Orlow and Wolf proposed the framework of HL and health outcomes, and hypothesized that SE might link HL and QOL [17]. Empirical studies have found that HL is associated with QOL via SE in patients with chronic disease, such as hypertension [18] or diabetes mellitus [19]. Up to now, much research has been done on the relationship between QOL and its related variables through correlation or regression method, which could only reflect the direct effect on QOL, and little work has revealed the pathway between HL, SE and QOL in CHD patients. Therefore, the present research was to investigate the pathway among HL, SE and QOL, and to investigate direct and indirect effects among variables in CHD patients. Significant associations were hypothesized as follows: (1) HL was associated with QOL directly in CHD patients; (2) SE was the mediator between HL and QOL in CHD patients. The study

may be used as the basis for developing interventions for improving the QOL in CHD patients.

Methods

Research designs and setting

A cross-sectional design was employed in the study. Patients who visited the outpatient clinic for post-discharge follow-up were selected through a convenience sampling following the STROBE guidelines from three tertiary general hospitals (the affiliated hospital of Hebei University, the first central hospital of Baoding, and the second hospital of Baoding) in Baoding City, Hebei province, mainland China. These hospitals are public, not referral, and each with >500 beds. Meanwhile, they are teaching hospitals of Hebei University. The research was approved by the Institutional Review Board of Hebei University. All methods were performed in accordance with the Declaration of Helsinki.

Participants

Participants who were selected for the study met the following criteria: (1) at least 18 years old; (2) diagnosis with CHD by a cardiologist (i.e., patients with myocardial infarction and/or PCI, angina) for more than 6 months; (3) able to give written informed consent. Participants who couldn't communicate independently because of cognitive or mental impairment or had severe complications (i.e., class IV heart failure, renal failure, diabetes, malignant tumor) were excluded. Each patient could join the study only once.

The sample size formula is $N = (u_{\alpha/2} \times \sigma / \delta)^2$. We expected to observe HL level. After the preliminary experiment showed that the standard deviation of HL was 4.1, based on the power of 80% and α of 5%, therefore, the sample size was 101 $[(1.96 * 4.1 / 0.8)^2 = 101]$. After 10% dropout rate was considered, the minimum estimation of final sample size was 111. Meanwhile, the sample included in structural equation modelling is minimum of 10 per indicator. The number of parameters to be estimated in this study was 17. As such, the minimum sample size is 170. A convenient sample of 200 patients could meet the sample size.

Data collection and measures

The data were collected from December 2018 to June 2019. Patients who met the criteria and visited the outpatient clinic were viewed as potential participants. They were informed the study and invited to participate by the researchers. If they agreed, they would be required to sign a consent. After informed consent was obtained, each participant was given the questionnaire. Patients could finish the questionnaire independently or with the

researchers' help in 20–30 min in a quiet room in the outpatient clinic.

Socio-demographic data

Gender, age, marital status, education attainment, monthly household income, employment status, family history of CHD and PCI treatment were included.

Health literacy scale

HL was assessed using self-reported Brief Screening Questions, which was used in patients with CVD and other diseases [14, 20–22]. There were 3 items as follows: “how confident are you filling out medical forms?”; “how often do you have problems learning about your medical condition because of difficulty understanding writing information?”; and “how often do you have someone help you read hospital materials?”. Answers were based on a 5-point Likert scale (1–5). The scores ranged from 3 to 15, with lower scores reflecting higher HL. Cronbach’s α was 0.88 in this study indicating excellent internal consistency.

Self efficacy scale

SE was assessed using a scale for chronic disease, which was used in some studies [23]. It included 2 dimensions and 6 items. The scores ranged from 6 to 60, and the higher the score, the higher the subjects' SE. Cronbach’s α was 0.94 in this study indicating excellent internal consistency.

Health-related quality of life

The Chinese version of the 36-item Short Form was used to assess the QOL [24]. It included 8 dimensions, and could be viewed as 2 summaries: physical health and mental health. The raw scores of each dimension were converted to a range 0 to 100. The higher score gets, the better level of QOL is. Cronbach’s α was 0.89 in our study indicating excellent internal consistency.

Ethical consideration

The research was approved by the Institutional Review Board of Hebei University, China. All participants could refuse or withdraw from the research at any time with no penalty.

Data analysis

All data were analysed using SPSS, version 22.0 and AMOS, version 24.0. Types and prevalence of demographic data were described with frequency and percentages. Total HL, SE and QOL scores were described by means and standard deviation. The questionnaire would be excluded for absence of more than 3 data. The independent sample *t* test and ANOVA test were applied to

analyze the difference in demographic characteristics of HL, SE and QOL. Pearson correlation was used to analyze the correlation between HL, SE and QOL. *P* values less than 0.05 indicated a significant difference.

The variables of HL, SE, and QOL were considered as latent constructs. The corresponding variables were considered as observed variables for each latent constructs. Since HL was measured in reverse, our assumptions were as follows: significant negative associations were observed from HL to QOL, from HL to SE, and possible associations were observed from SE to QOL. A structure equation model (SEM) was used to construct and validate the pathways. Comparative fit index (CFI) value above 0.9, root mean square error of approximation (RMSEA) value below 0.07, goodness-of-fit index (GFI) value above 0.9, adjust goodness-of-fit index (AGFI) value above 0.9, and the value of Chi-square/degrees of freedom (χ^2/df) below 3 indicated model fit.

Results

Distribution of related variables and associations among them

Of 232 patients invited, 22 patients were excluded for not meeting the inclusion criteria. 10 patients were excluded for missing data. Ultimately, 200 patients completed the questionnaires (Fig. 1). Table 1 shows the distribution of personal characteristics and their associations with the research valuables. We found that there were significant differences in HL and QOL among people of different ages and genders. Also, patients with different levels of education, employment status and monthly incomes had




Fig. 1 Sampling frame

Table 1 Personal characteristics and their associations with health literacy, self efficacy, and quality of life (n = 200)

Personal characteristics	Mean (SD) /n (%)	Health literacy Mean (SD)	Self efficacy Mean (SD)	Quality of life Mean (SD)
Age	65.51 (10.37)	9.6 (3.5)	28.8 (13.9)	381.8 (130.1)
r value		0.379**	-0.121	-0.187**
Gender				
Male	111 (55.5%)	8.2 (3.1)	30.1 (14.5)	402.9 (122.9)
Female	89 (44.5%)	11.3 (3.2)	27.2 (12.9)	355.7 (134.7)
t value		-6.9***	1.4	2.6*
Marital status				
Single(includes divorced, widowed, separated)	13 (6.5%)	10.7 (3.4)	31.1 (13.7)	382.0 (108.3)
Married	187 (93.5%)	9.5 (3.5)	28.7 (13.9)	381.8 (131.7)
t value		1.2	0.6	0.005
Education attainment				
Less than junior high school	166 (83.0%)	10.3 (3.2)	27.6 (13.4)	367.0 (123.8)
Senior high school/college/equivalent	34 (17.0%)	6.2 (2.7)	34.6 (14.7)	454.2 (137.5)
t value		6.9***	-2.7**	-3.7***
Income status (RMB/month)				
Below 2000	65 (32.5%)	11.1 (3.0)	27.1 (12.3)	337.2 (116.7)
2000-3000	81 (40.5%)	9.5 (3.1)	28.3 (13.0)	381.8 (116.1)
Above 3000	54 (27.0%)	8.0 (3.9)	31.7 (16.4)	435.6 (146.2)
F value		12.23***	1.8	9.1***
Employment status				
Unemployed	81 (40.5%)	10.8 (3.3)	25.8 (12.5)	358.9 (122.0)
Retired	92 (46.0%)	9.6 (3.2)	29.6 (13.3)	373.4 (121.4)
Employed	27 (13.5%)	6.1 (2.7)	35.2 (17.0)	479.4 (142.8)
F value		22.9***	5.2**	9.9***
Family history of CHD				
Yes	62 (31.0%)	9.4 (3.8)	29.7 (15.1)	391.8 (141.9)
No	138 (69.0%)	9.7 (3.3)	28.4 (13.3)	377.4 (124.7)
t value		-0.5	0.6	0.7
PCI treatment				
Yes	99 (49.5%)	9.5 (3.5)	29.8 (12.8)	389.4 (119.7)
No	101 (50.5%)	9.7 (3.5)	27.9 (14.9)	374.5 (139.7)
t value		-0.4	0.9	0.8

*p < 0.05

**p < 0.01

***p < 0.001

different HL, SE and QOL (see details in Table 1). Table 2 shows the associations among HL, SE and QOL. Significant associations were also found between them (see details in Table 2).

Test of the model

The significant direct pathways were found from HL to QOL, and SE played a partial mediating role between HL and QOL in the CHD population (see details in Fig. 2).

Table 2 Correlation among health literacy, self efficacy, and quality of life (n = 200)

	Health literacy	Self efficacy	Quality of life
Health literacy	1.000	-0.322***	-0.481***
Self efficacy		1.000	0.734***
Quality of life			1.000

***p < 0.001




Fig. 2 Final model and standardized pathway coefficients among health literacy, self efficacy, and quality of life. HL = health literacy; SE = self efficacy; QOL = quality of life; X1 = help read; X2 = confidence; X3 = problem learning; X4 = symptom management self efficacy; X5 = disease generic management self efficacy; Y1 = physical health; Y2 = mental health

The fit indices were $\chi^2/df=0.966$, $GFI=0.986$, $AGFI=0.963$, $CFI=1.000$, and $RMSEA=0.000$ [90% confidence interval (0.000–0.072)]. It indicated that the proposed model was acceptable. HL significantly indirectly affected QOL through SE. The direct effect of HL on SE was -0.32 , the direct effect of SE on QOL was 0.65 , the direct effect of HL on QOL was -0.27 , the indirect effect of HL on QOL was -0.21 (-0.32×0.65), and the total effect of HL on QOL was -0.48 ($-0.27 - 0.21$). Therefore, the total effect was greater than the direct effect, the direct effect was greater than the indirect effect of HL on QOL, and SE only played a partial mediating role. Lastly, HL accounted for 10.4% of the total variance in SE. HL and SE explained for 59.6% of the total variance in QOL.

Discussion

Several studies explored the relationship among HL, SE and QOL in patients with chronic disease [15, 25, 26]. However, correlation or regression method were often

used, which could only reflect the direct effect on QOL. Up to now, little work has revealed the indirect effect among HL, SE and QOL in CHD patients. Therefore, current study constructed and validated the pathways between HL, SE and QOL in CHD patients by SEM.

HL was significantly correlated with QOL in CHD patients in this study, which is in line with previous findings [13, 14], suggesting that CHD patients with inadequate HL tend to have poor QOL. Patients with chronic diseases often require medication therapy and changes in lifestyle for a long time [27]. Inadequate HL was related with less understanding of medical conditions [28]. Therefore, inadequate HL may affect health status. The results reported by Son et al. [14] revealed that adequate HL is a contributing factor in improving QOL in patients after PCI.

SE was the mediator between HL and QOL in CHD patients. SE directly positively influenced QOL in CHD patients. The finding was consistent with previous findings [29, 30], suggesting that CHD patients with low SE

tend to have poor QOL. Patients with higher SE may put in more effort, and have a great tendency in persisting in their attempts [31]. Also, our study confirmed that SE was the mediator between HL and QOL, revealing that CHD patients with inadequate HL have a lower SE in daily life, and ultimately leading to poor QOL, which echoes the previous framework proposed by Paasche-Orlow et al. [17].

Up to now, literature on HL in CHD patients is limited and the level of HL in these patients is generally low, accompanied by low QOL [32]. This study provides a model for describing the relationship among HL, SE and QOL in CHD patients. Health care givers can design intervention programs for CHD patients according to this model.

There are several limitations in the study. First, the pathways among HL, SE and QOL were from a cross-sectional study database. Accordingly, causal pathways in the model need to be confirmed through longitudinal and experimental studies. Second, a convenience sampling was used. Therefore, these findings cannot be extrapolated to CHD patients in other countries. Third, there are other modifiable factors that may influence QOL, such as depression, anxiety and social support [33], body mass index [34] and waist circumference [35]. Future research could expand the previous framework with the consideration of these variables. Fourth, the HL measurement in this study was functional, which didn't include communicable and critical literacy, thus, may not be perfect. Nevertheless, functional HL is the most commonly used measurement in mainland China [36]. In future studies, these factors should be considered.

Conclusions

In this investigation, the aim was to assess the relationship among HL, SE and QOL in CHD patients, and to investigate direct and indirect effects among variables. SE directly positively influenced QOL values and HL and SE accounted for 59.6% of the total variance. HL not only directly affects QOL, but also indirectly affects QOL via SE. Health care givers can manage CHD patients based on the findings of our study. Because SE directly influences QOL, modifying SE is essential for improving QOL in CHD patients. To improve SE, a potential strategy is to enhance health literacy.

Abbreviations

AGFI: Adjust goodness-of-fit index; CFI: Comparative fit index; CHD: Coronary heart disease; CVD: Cardiovascular disease; GFI: Goodness-of-fit index; HL: Health literacy; PCI: Percutaneous coronary intervention; QOL: Quality of life; RMSEA: Root mean square error of approximation; SE: Self efficacy; SEM: Structure equation model; χ^2/df : Chi-square/degrees of freedom.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12872-022-02720-8>.

Additional file 1: Raw data SPSS.

Additional file 2: Raw data Excel.

Acknowledgements

Not applicable

Authors contributions

Yan Wang and Shaoying Du conceptualized the idea for this study and obtained research funding. Shaoying Du, Zhimin Feng and Wen Wang contributed to the interview of the participant. Zhimin Feng, Wen Wang and Licong Tian contributed to the data analysis. Yan Wang is the guarantor and Shaoying Du drafted the manuscript. Zhimin Feng, Wen Wang and Licong Tian revised the manuscript. All authors read and approved the final manuscript.

Funding

Science and Technology Research Project of Hebei Higher Education Institutions QN2019178.

Availability of data and materials

All data generated or analysed during this study are included in this published article [and its Additional files 1 and 2].

Declarations

Ethical approval and consent to participate

The research was approved by the Institutional Review Board of Hebei University and was performed in accordance with the Declaration of Helsinki. All participants provided written informed consent.

Consent for publication

Not applicable.

Competing interests

Not applicable.

Author details

¹School of Nursing, Hebei University, No. 342 Yuhudong Road, Baoding City 071002, Hebei Province, People's Republic of China. ²School of Clinical Medicine, Hebei University, Baoding City, Hebei, People's Republic of China.

Received: 14 January 2022 Accepted: 13 June 2022

Published online: 23 June 2022

References

- Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, et al. Heart disease and stroke statistics-2017 update: a report from the American Heart Association. *Circulation*. 2017;135:e146-603.
- Hu S, Gao R, Liu L, Zhu M, Wang W, Wang Y, et al. Summary of the 2018 Report on cardiovascular disease in China. *Chin Circ J*. 2019;34(3):209–20 (**In Chinese**).
- Miguel RS, López-González AM, Sanchez-Iriso E, Mar J, Cabasés JM. Measuring health-related quality of life in drug clinical trials: is it given due importance? *Pharm World Sci*. 2008;30(2):154–60.
- Muhammad I, He HG, Kowitlawakul Y, Wang W. Narrative review of health related quality of life and its predictors among patients with coronary heart disease. *Int J Nurs Pract*. 2016;22(1):4–14.
- Muhammad I, He HG, Koh K, Thompson DR, Kowitlawakul Y, Wang W. Health-related quality of life and its predictors among outpatients with coronary heart disease in Singapore. *Appl Nurs Res*. 2014;27(3):175–80.
- Cassar S, Baldacchino DR. Quality of life after percutaneous coronary intervention: Part 2. *Br J Nurs*. 2012;21:1125–30.

7. Jankowska-Polańska B, Uchmanowicz I, Dudek K, Łoboz-Grudzień K. Impact of depression on quality of life results in patients following acute coronary syndrome treated with percutaneous coronary intervention (PCI). *J Nurs Care*. 2013;3(6):1–7.
8. Veenstra M, Pettersen KI, Rollag A, Stavem K. Association of changes in health-related quality of life in coronary heart disease with coronary procedures and sociodemographic characteristics. *Health Qual Life Out*. 2004;2:1–8.
9. Xiong Q, Zhou Q, Liu Y, Cao L, Liu X, He W, et al. Assessing quality of life among patients with coronary heart disease. *J Clin Cardiol*. 2014;30:27–30 (**[in Chinese]**).
10. Nielsen-Bohlman L, Panzer AM, Kindig DA. Health literacy: a prescription to end confusion. Washington, DC: The National Academies Press; 2004.
11. Peltzer S, Hellstern M, Genske A, Jünger S, Woopen C, Albus C. Health literacy in persons at risk of and patients with coronary heart disease: a systematic review. *Soc Sci Med*. 2019;245: 112711.
12. Nie XQ, Li YH, Li CN, Wu J, Li L. The association between health literacy and self-rated health among residents of China aged 15–69 years. *Am J Prev Med*. 2021;60(4):569–78.
13. Gonzalez-Chica DA, Minisi Z, Avery J, Duszynski K, Doust J, Tideman P, et al. Effect of health literacy on quality of life amongst patients with ischaemic heart disease in Australian general practice. *PLoS ONE*. 2016;11:e0151079.
14. Son YJ, Yu H. Influence of health literacy on HRQoL in patients after PCI. *West J Nurs Res*. 2016;38:1611–26.
15. Liu WC, Wang J, Luo YT, Huang HH, Xiao MC, Zhao QH. Use of structural equation modeling to identify factors of quality of life in coronary heart disease patient. *J Nurs Sci*. 2019;34(12):20–3 (**[in Chinese]**).
16. Dellafiore F, Pittella F, Arrigoni C, Baroni I, Conte G, Pasquale CD, et al. A multi-phase study for the development of a self-efficacy measuring scale for ostomy care nursing management. *J Adv Nurs*. 2020;76:409–19.
17. Paasche-orlow MK, Wolf MS. The causal pathways linking health literacy to health outcomes. *Am J Heal Behav*. 2007;31:519–26.
18. Wang C, Lang J, Xuan L, Li X, Zhang L. The effect of health literacy and self-management efficacy on the health-related quality of life of hypertensive patients in a western rural area of China: A cross-sectional study. *Int J Equity Health*. 2017;16(1):58.
19. Lee EH, Lee YW, Moon SH. A structural equation model linking health literacy to self-efficacy, self-care activities, and health-related quality of life in patients with type 2 diabetes. *Asian Nurs Res*. 2016;10:82–7.
20. McManus DD, Saczynski JS, Lessard D, Waring ME, Allison J, Parish DC, et al. Reliability of predicting early hospital readmission after discharge for an acute coronary syndrome using claim-based data. *Am J Cardiol*. 2016;117:501–7.
21. Peterson PN, Shetterly SM, Clarke CL, Bekelman DB, Chan PS, Allen LA, et al. Health literacy and outcomes among patients with heart failure. *J Am Med Assoc*. 2011;305:1695–701.
22. Chew LD, Bradley KA, Boyko EJ. Brief questions to identify patients with inadequate health literacy. *Fam Med*. 2004;36:588–94.
23. Fan X, Lv F. Psychosocial factors associated with self-efficacy for managing chronic disease in patients with chronic heart failure. *Eur J Cardiovasc Nur*. 2016;15(4):255–61.
24. Peng Z, Li H, Zhang C, Qian X, Feng Z, Zhu S. A retrospective study of chronic post-surgical pain following thoracic surgery: prevalence, risk factors, incidence of neuropathic component, and impact on quality of life. *PLoS ONE*. 2014;9(2): e90014.
25. Zhang J, Gilmour S, Liu Y, Ota E. Effect of health literacy on quality of life among patients with chronic heart failure in China. *Qual Life Res*. 2020;29:453–61.
26. Magon A, Caruso R, Sironi A, Mirabella S, Dellafiore F, Arrigoni C, et al. Trajectories of health-related quality of life, health literacy, and self-efficacy in curatively-treated patients with esophageal cancer: a longitudinal single-center study in Italy. *J Patient Exp*. 2021;8:1–10.
27. Smith SC, Allen J, Blair SN, et al. AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update: endorsed by the National Heart, Lung, and Blood Institute. *J Am Coll Cardiol*. 2006;47(10):2130–9.
28. Ownby RL, Acevedo A, Jacobs RJ, Caballero J, Waldrop-Valverde D. Quality of life, health status, and health service utilization related to a new measure of health literacy: FLIGHT/VIDAS. *Patient Educ Couns*. 2014;96:404–10.
29. Li C, Jiang F, Liu B. Relationship between quality of life and self-efficacy in coronary heart disease patients. *Chin J Nurs Sci*. 2009;24(11):36–7 (**[in Chinese]**).
30. Sarkar U, Ali S, Whooley MA. Self-efficacy and health status in patients with coronary heart disease: findings from the heart and soul study. *Psychosom Med*. 2007;69:306–12.
31. Polsook R, Aunguroch Y, Thongvichean T. The effect of self-efficacy enhancement program on medication adherence among post-acute myocardial infarction. *Appl Nurs Res*. 2016;32:67–72.
32. Ghisi GLM, Chaves GSDS, Britto RR, Oh P. Health literacy and coronary artery disease: a systematic review. *Patient Educ Couns*. 2017;101(2):177–84.
33. Nekouei ZK, Yousefy A, Doost HTN, Manshaee G, Sadeghei M. Structural model of psychological risk and protective factors affecting on quality of life in patients with coronary heart disease: a psychocardiology model. *J Res Med Sci*. 2014;19:90–8.
34. Hokkanen M, Järvinen O, Huhtala H, Laurikka J. The effect of obesity on long-term survival and health-related quality of life after coronary artery bypass grafting: a 12-year follow-up. *Coro Artery Dis*. 2018;29(5):378–83.
35. Hyun YY, Lee KB, Chung W, Kim YS, Han SH, Oh YK, et al. Body Mass Index, waist circumference, and health-related quality of life in adults with chronic kidney disease. *Qual Life Res*. 2019;28(4):1075–83.
36. Wang P, Mao Q, Tao M, Tian X, Li Y, Qian L, et al. Survey on the status of health literacy of Chinese residents in 2008. *Chin J Health Educ*. 2010;26:243–6 (**[in Chinese]**).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

