CORRECTION Open Access



Correction to: does stress perfusion imaging improve the diagnostic accuracy of late gadolinium enhanced cardiac magnetic resonance for establishing the etiology of heart failure?

Gaurav S. Gulsin^{1,2*}, Abhishek Shetye^{1,2}, Jeffrey Khoo^{1,2}, Daniel J. Swarbrick^{1,2}, Eylem Levelt^{1,2}, Florence Y. Lai^{1,2}, lain B. Squire^{1,2}, Jayanth R. Arnold^{1,2} and Gerry P. McCann^{1,2}

Correction to: BMC Cardiovascular Disorders (2017) 17:98. DOI: 10.1186/s12872-017-0529-y

Following publication of the original article [1], the author reported his name has erroneously spelled as Abishek Shetye. The correct name is Abhishek Shetye.

The publisher apologizes for any inconvenience caused by this error.

Received: 14 January 2019 Accepted: 14 January 2019 Published online: 21 January 2019

Reference

 Gulsin, et al. BMC Cardiovasc Disord. 2017;17:98. https://doi.org/10.1186/ s12872-017-0529-y.

²The NIHR Leicester Cardiovascular Biomedical Research Unit, Glenfield Hospital, Leicester, UK



^{*} Correspondence: gg149@leicester.ac.uk

¹Department of Cardiovascular Sciences, University of Leicester, Glenfield Hospital, Groby Road, Leicester LE3 9QP, UK